

# OpenType math font Fira

Herbert Voß

January 16, 2023

## Contents

<b>1</b>	<b>Usage</b>	<b>1</b>
<b>2</b>	<b>The default regular weight</b>	<b>2</b>
2.1	Version normal . . . . .	2
2.2	Version bold . . . . .	2
<b>3</b>	<b>Examples</b>	<b>2</b>
3.1	Digits . . . . .	2
3.2	Alphabets . . . . .	3
3.3	Equations test . . . . .	4

### Abstract

The math font FIRA is derived from the Fira Sans and Fira Go sans serif. There are several math versions available (<https://github.com/Stone-Zeng/FiraMath/>) but only the regular version has from todays update all symbols.

## 1 Usage

\usepackage[<options>]{firamath-otf}

Optional arguments are

**fakebold** Use faked bold symbols

**usefilenames** Use filenames for the fonts instead of the symbolic font names

All other unknown options, e.g. `mathrm=sym` will be passed to the main package `unicode-math`.

The package itself loads by default

```
\RequirePackage{iftex,xkeyval,textcomp}  
\RequirePackage{unicode-math}
```

## 2 The default regular weight

### 2.1 Version normal

$$\begin{aligned}\frac{\partial \rho}{\partial t} + \operatorname{div}(\rho \vec{v}) &= 0 \\ \rho \frac{\partial \vec{v}}{\partial t} + (\rho \vec{v} \cdot \nabla) \vec{v} &= \vec{f}_0 + \operatorname{div} T = \vec{f}_0 - \operatorname{grad} p + \operatorname{div} T' \\ \rho T \frac{ds}{dt} &= \rho \frac{de}{dt} - \frac{p}{\rho} \frac{d\rho}{dt} = -\operatorname{div} \vec{q} + T' : D\end{aligned}\quad (1)$$

$$\frac{\partial}{\partial t} \iiint \rho d^3V + \oint \rho (\vec{v} \cdot \vec{n}) d^2A = 0 \quad (2)$$

$$\frac{\partial}{\partial t} \iiint \rho \vec{v} d^3V + \oint \rho \vec{v} (\vec{v} \cdot \vec{n}) d^2A = \iiint f_0 d^3V + \oint \vec{n} \cdot T d^2A \quad (3)$$

$$\begin{aligned}\frac{\partial}{\partial t} \iiint \left( \frac{1}{2} v^2 + e \right) \rho d^3V + \oint \left( \frac{1}{2} v^2 + e \right) \rho (\vec{v} \cdot \vec{n}) d^2A &= \\ - \oint (\vec{q} \cdot \vec{v} \vec{n}) d^2A + \iiint (\vec{v} \cdot \vec{f}_0) d^3V + \oint (\vec{v} \cdot \vec{n} T) d^2A.\end{aligned}\quad (4)$$

### 2.2 Version bold

The bold characters are created with the optional argument `fakebold` which loads the package `xfakebold` which writes some information into the created PDF to get bold characters. For more informations see the documentation of `xfakebold`.

$$\frac{\partial}{\partial t} \iiint \rho d^3V + \oint \rho (\vec{v} \cdot \vec{vecn}) d^2A = 0 \quad (5)$$

$$\frac{\partial}{\partial t} \iiint \rho \vec{v} d^3V + \oint \rho \vec{v} (\vec{v} \cdot \vec{n}) d^2A = \iiint f_0 d^3V + \oint \vec{n} \cdot T d^2A \quad (6)$$

$$\begin{aligned}\frac{\partial}{\partial t} \iiint \left( \frac{1}{2} v^2 + e \right) \rho d^3V + \oint \left( \frac{1}{2} v^2 + e \right) \rho (\vec{v} \cdot \vec{n}) d^2A &= \\ - \oint (\vec{q} \cdot \vec{vecn}) d^2A + \iiint (\vec{v} \cdot \vec{f}_0) d^3V + \oint (\vec{v} \cdot \vec{n} T) d^2A.\end{aligned}\quad (7)$$

## 3 Examples

### 3.1 Digits

- Digits:

0123456789

- Proportional digits: 0123456789
- Bold digits (`\symbf`): **0123456789**
- Bold proportional digits (`\symbf`): **0123456789**

### 3.2 Alphabets

- Latin letters (mathnormal):  
 $ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz$
- Latin upright letters (`\symup`):  
 $ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz$
- Latin typewriter letters (`\syttt`):  
 $ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz$
- Latin bold letters (`\symbf`):  
 **$ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz$**
- Latin bold upright letters (`\symbfup`):  
 **$ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz$**
- Latin blackboard letters (`\symbb`):  
 $ABCDEFGHIJKLMNOPQRSTUVWXYZabcdefghijklmnopqrstuvwxyz$
- Greek letters:  
 $ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεεζηθικιλμνξοπρσςτυφχψω$
- Greek upright letters (`\symup`):  
 $ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεεζηθικιλμνξοπρσςτυφχψω$
- Greek bold letters (`\symbf`):  
 **$ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεεζηθικιλμνξοπρσςτυφχψω$**
- Greek bold upright letters (`\symbfup`):  
 **$ΑΒΓΔΕΖΗΘΙΚΛΜΝΞΟΠΡΣΤΥΦΧΨΩαβγδεεζηθικιλμνξοπρσςτυφχψω$**
- Dotless letters:  
 $I + J + I + J$
- Hebrew נ + ג + ג + ת
- Ligature (text):  
 $ff \ fi \ fl \ ffi \ ffl$
- Non-ligature (math):  
 $ff \ fi \ fl \ ffi \ ffl + ff \ fi \ fl \ ffi \ ffl + ff \ fi \ fl \ ffi \ ffl$
- Miscellaneous:  
 $\ddot{h} + \ddot{h} + \ddot{A}$   
 $\forall x > x_0, \exists \delta, \delta \in \emptyset$

### 3.3 Equations test

- Basic:

$$1 + 2 - 3 \times 4 \div 5 \pm 6 \mp 7 \div 8 = -a \oplus b \otimes c$$

- Binary relations  $x + - \oplus \otimes \ominus \odot \odot \cdots \times \div y$

- Set theory  $A \cap B \cup C \sqcap D \sqcup R \sqcup k \sqcup l \sqcup m$

$$A \subset B \supset C \subseteq D \supseteq E \quad F \supset G + A \sqsubset B \sqsupset C \sqsubseteq D \supseteq E$$

$$\mathcal{C}_U A \cup \mathcal{C}_C C \subset \mathcal{C}_U A \cup \mathcal{C}_C C \in R \in Q \ni Z \ni N$$

- Superscript and subscript:

$$2^2 + 2^{2^2} + 2^{2^{2^2}} + 2^{2^{2^2}} + x_a + x_{a_i} + x_{a_{i_1}}$$

- Arrows:

$$x \leftarrow y \rightarrow z \leftrightarrow w \Leftarrow y \Rightarrow z \Leftrightarrow w \Leftarrow a \Rightarrow b \Leftrightarrow c \Leftarrow a \Rightarrow b \Leftrightarrow c$$

$$x \uparrow y \downarrow z \Updownarrow w \uparrow a \Downarrow b \Updownarrow c$$

$$p \nwarrow p \nearrow p \searrow p \swarrow p \nearrow p \searrow p \swarrow p$$

$$x \leftarrow x \leftarrow x \uparrow x \rightarrow x \rightarrow x \downarrow x \downarrow x$$

$$A \leftarrow B \rightarrow C \leftrightarrow D \Leftarrow E \Rightarrow F \Leftrightarrow G$$

$$X \leftrightarrow Y \mapsto Z \uparrow W \downarrow P \Leftarrow S \Rightarrow R$$

$$M \leftarrow N \mapsto O \Leftarrow K \Rightarrow L$$

$$f \rightleftarrows f \uparrow\!\!\!l f \leftrightharpoons f \downarrow\!\!\!l g \Rightarrow g \uparrow\!\!\!l g \Leftarrow g \Downarrow h \Rrightarrow h \Lleftarrow p \Leftarrow p \Rightarrow p \Downarrow\!\!\!l p \Downarrow\!\!\!l p$$

- Math accents:

$$\acute{x} \check{x} \ddot{x} \hat{x} \ddot{\check{x}} \acute{\check{x}} \acute{x} \check{\acute{x}} \ddot{\acute{x}} \check{\acute{x}} \acute{\check{x}} \acute{\acute{x}} \acute{\check{\acute{x}}} \acute{\acute{\check{x}}} \acute{\acute{\acute{x}}} \acute{\acute{\acute{\check{x}}}} \acute{\acute{\acute{\acute{x}}}} \acute{\acute{\acute{\acute{\check{x}}}}}$$

- Integral:

$$\int_0^\pi \sin x \, dx = \int_0^\pi \sin x \, dx = \cos 0 - \cos \pi = 2$$

$$\int_{-\infty}^{+\infty} dz \iint_{-\infty}^{+\infty} d^2y \iiint_{-\infty}^{+\infty} d^3x \iiint_{-\infty}^{+\infty} d^4p$$

$$\oint dr \iint d\theta \iiint d\varphi$$

$$\int_0^\pi \sin x \, dx = \int_0^\pi \sin x \, dx = \cos 0 - \cos \pi + C$$

$$\int_{-\infty}^{+\infty} dz \iint_{-\infty}^{+\infty} d^2y \iiint_{-\infty}^{+\infty} d^3x \iiint_{-\infty}^{+\infty} d^4p$$

$$\oint dr \iint d\theta \iiint d\varphi$$

- Huge operators:

$$\int_0^\infty \int_0^\infty \sum_{i=1}^\infty \prod_{j=i}^\infty \prod_{k=i}^\infty$$

$$\sum_{i=1}^\infty \frac{1}{x^i} = \frac{1}{1-x} \quad \prod_{i=1}^\infty \frac{1}{x^i} = x^{-n(n+1)/2} \quad \prod_{i=i}^\infty \frac{1}{x^i} = ?$$

- Huge operators (inline):

$$\int_0^\infty \int_0^\infty \iint dx \iiint dy \iiii dp \oint dr \oint\oint d\theta \oint\oint\oint d\varphi \sum_{i=1}^\infty \prod_{j=i}^\infty \prod_{i=i}^\infty$$

- Huge operators (inline):

$$\int_0^\infty \int_0^\infty \iint dx \iiii dy \iiii dp \oint dr \oint\oint d\theta \oint\oint\oint d\varphi \sum_{i=1}^\infty \prod_{j=i}^\infty \prod_{i=i}^\infty$$

- Fraction:

$$\frac{1}{2} + \frac{1}{\frac{2}{3} + 4} + \frac{\frac{1}{2} + 3}{4}$$

- Fraction (inline):

$$\frac{1}{2} + \frac{1g}{2} + \frac{1}{\frac{2}{3} + 4} + \frac{\frac{1}{2} + 3}{4}$$


- Radical:

$$\sqrt{2} + \sqrt{2^2} + \sqrt{1 + \sqrt{2}} + \sqrt{1 + \sqrt{1 + \sqrt{3}}} + \sqrt{\sqrt{\sqrt{2}} + \sqrt{\frac{1}{2}}}$$

$$\sqrt[3]{2} + \sqrt[3]{2^2} + \sqrt[3]{1 + \sqrt[3]{2}} + \sqrt[3]{1 + \sqrt[3]{1 + \sqrt[3]{3}}} + \sqrt[3]{\sqrt[3]{\sqrt[3]{2}}} + \sqrt[3]{\frac{1}{2}}$$

$$\sqrt[4]{2} + \sqrt[4]{2^2} + \sqrt[4]{1 + \sqrt[4]{2}} + \sqrt[4]{1 + \sqrt[4]{1 + \sqrt[4]{3}}} + \sqrt[4]{\sqrt[4]{\sqrt[4]{2}}} + \sqrt[4]{\frac{1}{2}}$$

$$\sqrt[x]{y} + \sqrt[x]{\sqrt[y]{y}} + \sqrt[x]{\sqrt[x]{\sqrt[y]{y}}} + \sqrt[x]{\frac{1}{2}} + \sqrt[x]{\frac{x}{z} + \sqrt[x]{\frac{y}{w}}} + \sqrt[x]{\sqrt[x]{\frac{y}{z}} + \sqrt[x]{\sqrt[w]{\frac{y}{p}}}} + \sqrt[x]{\sqrt[x]{\frac{y}{z}} + \sqrt[x]{\sqrt[w]{\frac{y}{p}}}} + \sqrt[x]{\sqrt[x]{\frac{y}{z}} + \sqrt[x]{\sqrt[w]{\frac{y}{p}}}}$$



- Brackets:

$$\begin{aligned}
 & (a)(A)(O)(Y)(y)(f)(Q)(T)(Y)(j)(q) \\
 & \left( \left( \left( (x) \right) \right) \right) \quad \left( \left( \left( (x) \right) \right) \right) \quad \left[ \left[ \left[ [x] \right] \right] \right] \quad \left\{ \left\{ \left\{ \{x\} \right\} \right\} \right\} \\
 & (x) + (x^2) + \left( \frac{1}{2} \right) + \left( \frac{2^2}{3} \right) + \left( \frac{\frac{1}{2}}{\frac{3}{4}} \right) \\
 & (\{\})[\{\}] \{ \{\} \} \quad \left( \left( \right) \right) \left[ \left[ \right] \right] \left\{ \left\{ \right\} \right\} \quad \left( \left( \right) \right) \left[ \left[ \right] \right] \left\{ \left\{ \right\} \right\}
 \end{aligned}$$

- More brackets:

[ceiling] [floor] (group)

- Bra-kets:

$$\langle x| + |x\rangle + \langle\alpha|\beta\rangle + |\alpha^2\rangle\langle\beta^2| + \left|\frac{1}{2}\right| + \left|\frac{1}{2}\right\rangle + \left\langle\frac{1}{2}\middle|\frac{1}{2}\right\rangle + \left|\frac{1}{2}\right\rangle\left\langle\frac{1}{2}\right| + \left\langle\frac{a^2}{b^2}\right| + \left|\frac{e^{x^2}}{e^{y^2}}\right\rangle$$

$$\langle \rangle \langle \rangle$$

- Matrices:

$$\begin{array}{c} \begin{pmatrix} a & b \\ c & d \end{pmatrix} + \begin{pmatrix} a & b \\ c & d \end{pmatrix} \\ \begin{pmatrix} a & b & c & d \\ x & y & z & w \end{pmatrix} \quad \begin{bmatrix} a & b & c & d \\ x & y & z & w \end{bmatrix} \quad \left\{ \begin{matrix} a & b & c & d \\ x & y & z & w \end{matrix} \right\} \quad \left| \begin{matrix} a & b & c & d \\ x & y & z & w \end{matrix} \right| \quad \left\| \begin{matrix} a & b & c & d \\ x & y & z & w \end{matrix} \right\| \\ \begin{pmatrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{pmatrix} \quad \begin{bmatrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{bmatrix} \quad \left\{ \begin{matrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{matrix} \right\} \quad \left| \begin{matrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{matrix} \right| \quad \left\| \begin{matrix} a & b & c & d \\ k & l & m & n \\ x & y & z & w \end{matrix} \right\| \\ \begin{pmatrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{pmatrix} \quad \begin{bmatrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{bmatrix} \quad \left\{ \begin{matrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{matrix} \right\} \quad \left| \begin{matrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{matrix} \right| \quad \left\| \begin{matrix} a & b & c & d \\ k & l & m & n \\ p & q & s & t \\ x & y & z & w \end{matrix} \right\| \end{array}$$

- Nablas:

$$\nabla x + \nabla f + \nabla \cdot \mathbf{u} + \nabla \times \mathbf{v}$$

$$\nabla \quad \nabla \quad \nabla \quad \nabla; \quad \tilde{\nabla} \quad \tilde{\nabla} \quad \tilde{\nabla} \quad \tilde{\nabla}$$

- Over-/underline and over-/underbraces

$$\begin{array}{ccccccccc} \overline{b} & \overline{ab} & \overline{abc} & \overline{abcd} & \overline{abcde} & \overline{a+b+c} & \overbrace{x_1, x_2, \dots, x_n} \\ \overbrace{\widehat{b}} & \widehat{\overline{ab}} & \widehat{\overline{abc}} & \widehat{\overline{abcd}} & \widehat{\overline{abcde}} & \widehat{a+b+c} & \overbrace{\widehat{x_1, x_2, \dots, x_n}}^n \\ \overbrace{\widehat{\overline{b}}} & \widehat{\overline{ab}} & \widehat{\overline{abc}} & \widehat{\overline{abcd}} & \widehat{\overline{abcde}} & \widehat{a+b+c} & \overbrace{\widehat{x_1, x_2, \dots, x_n}}^n \\ \overbrace{\widehat{\overline{b}}} & \widehat{\overline{ab}} & \widehat{\overline{abc}} & \widehat{\overline{abcd}} & \widehat{\overline{abcde}} & \widehat{a+b+c} & \overbrace{\widehat{x_1, x_2, \dots, x_n}}^n \\ \underline{b} & \underline{ab} & \underline{abc} & \underline{abcd} & \underline{abcde} & \underline{a+b+c} & \overbrace{x_1, x_2, \dots, x_n} \\ \underline{\overbrace{b}} & \underline{\overbrace{ab}} & \underline{\overbrace{abc}} & \underline{\overbrace{abcd}} & \underline{\overbrace{abcde}} & \underline{\overbrace{a+b+c}} & \overbrace{\underline{x_1, x_2, \dots, x_n}}^n \\ \underline{\overbrace{b}} & \underline{\overbrace{ab}} & \underline{\overbrace{abc}} & \underline{\overbrace{abcd}} & \underline{\overbrace{abcde}} & \underline{\overbrace{a+b+c}} & \overbrace{\underline{x_1, x_2, \dots, x_n}}^n \\ \underline{\overbrace{b}} & \underline{\overbrace{ab}} & \underline{\overbrace{abc}} & \underline{\overbrace{abcd}} & \underline{\overbrace{abcde}} & \underline{\overbrace{a+b+c}} & \overbrace{\underline{x_1, x_2, \dots, x_n}}^n \end{array}$$

- Primes

$$x' x'' x''' x'''' x^{x'} x^{x''} x^{x'''} x^{x''''} x^{x^{'}}$$

$$x' x'' x''' x''''$$

$$x' x'' x''' x''''$$

$$\lim_{x \rightarrow \infty} \frac{1}{x^2} = 0$$

$$\frac{\partial y(x)}{\partial x} = \frac{dy(x)}{dx} = y'(x)$$

$$\frac{\partial y(x)}{\partial x} = \frac{dy(x)}{dx} = y'(x)$$